SSWP 2018 Non-Native Invasive Plant (NNIP) Species Datasheet

Datasheet ID : 20180416-SSWP-NNIP-CENMEL-3

(Date-SSWP-Location-NNIP-SurveyTeam#- Datasheet#
e.g. 20180420-SSWP-PL-NNIP-PL-1-2)

Date	Location[1]	Survey Team # (1, 2, 3, or 4)	Survey Team Daily NNIP Datasheet#	Photo Location (iPad, field camera, etc)	USFS Land?
4-16-18	Castaic T-line	1	3	iPad	N

Recorder (First Initial & Last Name)	Habitat Description (habitat type and dominant species)
S. Tona & A. Sorci	Chamise CA buckwheat scrub

Land Use Notes (surrounding land use, visible disturbance, source of continued infestation, etc.)

Are target NNIP species too widespread or variable for mapping to remain accurate? Describe below
(Species Code, approximate extent, etc.)

N

Other (non-target) NNIP species present in this location

Brassica nigra, Avena barbata, Bromus hordeaceus

Variances/Other Notes (e.g.,

Target Species Data (refer to target species list)
(note, if not on USFS land, only record CDFA-listed species below)

Species Code	CDFA Status	C or D[2]	D or W[3]	Percent Cover[4] (Absolute)	Percent Phenology[4]			Estimated Area (sq.ft.)	Occurrence Extends Beyond the Project Boundary? (Y, N, or N/A)
					Vegetative	Flower	Fruiting		
CENMEL	C	C	D	2	100			1,090 sq ft	N

[1]Location = Quail Lake (QL), Gorman Creek (GC), Pyramid Lake (PL), Warne Transmission Line (WT), Castaic Creek (CC), Elderberry Forebay (EF), or Castaic Transmission Line (CT)
[2]C or D = Concentrated or Diffuse
[3]D or W – Discrete or Widespread (a GPS point can be taken, rather than polygon, if the occurrence is discrete)
[4]Percent cover classes: 1 = 0-1%, 2 = 2-5%, 3 = 6-10%, 4 =11-15%, 5 = 16-25%, 6 = 26-35%, 7 = 36-50%, 8 = 51-75%, 9 = 76-100%